



Make the right decision

Ready-to-use gel bait

Habitro Ant Gel is a ready-to-use insecticidal bait for the rapid control of a variety of ants

What is Habitro Ant Gel?

Habitro Ant Gel is a viscous liquid ant bait made of 99.9% naturally derived ingredients for the control of ants.

It's patented formulation (HVL Technology) naturally preserves the bait to ensure long term attractiveness without artificial preservatives.



Pharaoh's Ant

Ghost Ant





// Control sugar feeding species as well as certain protein feeding ant species.

Unique delayed-action kill that eliminates the entire colony

- // Ants take the bait back to the nest where the active ingredient kills the gueen, workers and the rest of the nest.
- // Significant reduction in ant activity in days, followed by colony elimination in weeks (depending on ant species & size of nest)

Utilises patented HVL Technology

// Extended bait life up to minimum 8 weeks after application.





A completely homogeneous viscous liquid bait

- // Clear formulation allows for inconspicuous bait placements.
- // Matches the natural preferences of the target pest.
- // Easy to apply

and









Who can use Habitro Ant

Indoor and outdoors of

buildings.

commercial and residential

How to use:

Habitro Ant Gel comes with a ready-to-use applicator syringe.



1. Open the cap of the svrinae.



2. The product is ready-touse.




Directions for use

For maximum effectiveness, Habitro Ant Gel must be placed on or near ant trails:

- Apply one spot of 0.2 g per linear meter of ant trail (equivalent to one pea-sized drop or a number of smaller drops)
- For outdoor use. Habitro Ant Gel can be placed in cracks and crevices (in brick walls, between pavement slabs, etc.) where ants are seen to be active or when ant trails are evident.
- If nest entrances can be identified, inject 2 g of product directly







Best practice guidelines for ant control

- Conduct a thorough inspection to get a full idea of extent of ant trails and possible nesting sites.
- Place baits close to foraging ant workers.
- Correct any conditions which may be contributing to the infestation (e.g. food spillages, honeydew producing insects on plants, accessible food etc.)
- Ensure an adequate volume of bait and/or bait placements relative to infestation or feeding levels.
- Place baits preferably in an area protected by the rain.
- It is recommended to monitor at day 3, 7, 14 and 28 days after treatment. Repeat treatment when infestations are high.



Bayer Co. (Malaysia) Sdn Bhd Reg. No. 196701000620 (7563-M) **Crop Science Division, Environmental Science** 25-03 & 25-04, Level 25, IMAZIUM No 8, Jalan SS21/37 Damansara Uptown 47400 Petaling Jaya, Selangor Malaysia